

Bismuth compounds in organic synthesis. Deprotection of ketoximes using bismuth bromide-bismuth triflate

Joshua N. Arnold, Patrick D. Hayes, Robert L. Kohaus and Ram S. Mohan

Abstract—Ketoximes undergo deprotection in CH₃CN/acetone/H₂O (3:6:1) in the presence of 20–40 mol% BiBr₃/5 mol% Bi(OTf)₃. Bismuth(III) salts are relatively non-toxic, insensitive to air and inexpensive. These features coupled with the use of a relatively non-toxic solvent system make this method an attractive alternative to existing routes for deprotection of ketoximes.

Keywords: bismuth and compounds; oximes; deprotection; environment-friendly chemistry.


Oximes are frequently used to protect carbonyl compounds and hence considerable attention has been given to develop methods for their deprotection.¹ Oximes can also be synthesized from non-carbonyl compounds and thus their conversion to carbonyl compounds constitutes a useful synthesis of the latter.² The classical method for deprotection of oximes viz. hydrolytic cleavage requires the use of strong mineral acids and often results in low yields due to the formation of polymeric by-products. Hence a number of oxidative methods have been developed for the cleavage of oximes. Some examples include Dess–Martin periodinane,³ PCC,⁴ TBHP,⁵ I₂/CH₃CN,⁶ manganese acetate/benzene,⁷ and potassium peroxymonopersulfate.⁸ However, many of these reagents or the solvents used are toxic, corrosive or difficult to handle, especially on a large scale. With increasing environmental concerns, it is imperative that new 'environment friendly' reagents be developed.⁹ One example of a mild method for deprotection of oximes uses DOWEX-50 resin.¹⁰ Recently, bismuth compounds have become attractive

candidates for use as reagents in organic synthesis because most bismuth compounds are relatively non-toxic, readily available at low cost and are fairly insensitive to small amounts of water.¹¹ Bismuth has an electron configuration of [Xe]4f¹⁴5d¹⁰6s²6p³. Due to the weak shielding of the 4f electrons (Lanthanide contraction), bismuth(III) compounds exhibit Lewis acidity. BiCl₃/microwave,¹² BiCl₃ in the presence of benzyltriphenylphosphonium peroxymonosulfate¹³ and Bi(NO₃)₃·5H₂O¹⁴ have been used for the deprotection of oximes. We now report the use of a new bismuth-based catalyst system (bismuth bromide/bismuth triflate) for the deprotection of ketoximes.

Our continued work with bismuth(III) reagents has led to the development of bismuth bromide/bismuth triflate as an efficient reagent for the conversion of ketoximes to ketones (Scheme 1, Table 1).

The reaction is carried out using 20–40 mol% BiBr₃ and 5 mol% Bi(OTf)₃·4H₂O as a co-catalyst. The use of an inert atmosphere is not required for these reactions. Several solvents were investigated during the course of this study. The best results were achieved using CH₃CN/acetone/H₂O (3:6:1, v/v/v). The results of this study are summarized in Table 1.

While detailed mechanistic studies were not carried out, a few points merit comment. While BiBr₃ alone was effective in catalyzing the deprotection, the addition of Bi(OTf)₃ accelerated the reaction considerably. The use of 5 mol% Bi(OTf)₃ alone (no BiBr₃) was almost ineffective in catalyzing the deprotection. However, when 40 mol% Bi(OTf)₃ was used alone, the deprotection did



Scheme 1.

Table 1. Deprotection of ketoximes using $\text{BiBr}_3\text{-Bi}(\text{OTf})_3\cdot 4\text{H}_2\text{O}$ in $\text{CH}_3\text{CN}/\text{acetone}/\text{H}_2\text{O}$

Entry	Oxime	Time, h	Product	Yield (%) ^{a,b}
1		2.5		68 (94)
2		2		77 (98)
3		22		76 (98) ^c
4		3		79 (96)
5		0.7 h		65 (95)
6		4		85 (96) ^c
7		1		79 (93)
8		1		64 (92)

^aYields are not optimized. Refers to yield of isolated product. .

^bNumbers in parentheses refer to percent purity of crude product as estimated by ¹H & ¹³C NMR spectroscopy and GC analysis. The remainder was oxime in all cases.

^cProduct was purified by recrystallization from hexane.

proceed smoothly. Since BiBr_3 is commercially available, we found it more practical to use BiBr_3 with $\text{Bi}(\text{OTf})_3$ as a co-catalyst.¹⁵ A suspension of the catalyst system in water is acidic (pH 2). Hence it is possible that the true catalyst is HBr and/or TfOH, released from hydrolysis of BiBr_3 and $\text{Bi}(\text{OTf})_3$. Indeed, the deprotection of 1-indanone oxime using 1.2 equivalents of HBr and 15 mol% TfOH proceeded smoothly, but the highly corrosive nature of HBr and TfOH detract from their use. The attempted deprotection of 1-indanone oxime using BiBr_3 and $\text{Bi}(\text{OTf})_3$ in the presence of proton-sponge[®](*N,N,N',N'*-tetramethyl-1,8-naphthalenediamine)¹⁶ was unsuccessful suggesting that the main role of the catalysts is to provide an acidic reaction medium.

The choice of solvents also merits comments. Acetone/ H_2O was found to be effective but the rate of deprotection was considerably slower in this solvent system. For example, the deprotection of 1-indanone oxime in acetone/ H_2O took over 12 h, while the addition of the polar coordinating solvent CH_3CN accelerated the reaction considerably (reaction time=3 h). Acetone oxime was found to be a product of the reaction. This is not surprising in light of the fact that oximes have

been synthesized from aldehydes and ketones by acid catalyzed transoximations using acetone oxime.¹⁷ This suggests that the role of acetone is to drive the equilibrium towards product formation. The deprotection of 1-indanone oxime in acetonitrile/water (6:4) was only 78% complete after 5 h, suggesting that acetone plays a crucial role in driving the reaction. Water is also necessary for the deprotection since the reaction was significantly slower in $\text{CH}_3\text{CN}/\text{acetone}$ as the solvent.

The deprotection works well with a variety of ketoximes. Aldoximes on the other hand proved much more resistant to the deprotection conditions. With benzaldehyde oxime, the reaction reached a 1:1 equilibrium mixture of benzaldehyde and the starting oxime within an hour. When the reaction was carried out with 40 mol% $\text{BiBr}_3/20$ mol% $\text{Bi}(\text{OTf})_3$, the reaction reached a 3.5:1 equilibrium mixture of benzaldehyde and the starting oxime in 1 h.

A representative procedure is given here: A solution 1-indanone oxime (0.500 g, 3.40 mmol) in $\text{CH}_3\text{CN}/\text{acetone}/\text{H}_2\text{O}$ (3:6:1, v/v/v) (5.00 mL) was stirred as BiBr_3 (0.609 g, 1.36 mmol), and $\text{Bi}(\text{OTf})_3\cdot 4\text{H}_2\text{O}$ (0.111 g,

0.170 mmol) were added. The mixture was stirred and heated at reflux. Reaction progress was followed by GC analysis. After 3 h, 10% aqueous acetic acid was added and the mixture was suction filtered and the solids were rinsed with ether (20 mL). The combined filtrates were concentrated on a rotary evaporator to remove the organic solvents. The residue was extracted with ether (2×40 mL). The organic layer was washed successively with 2 M NaOH (15 mL), H₂O (3×15 mL), saturated aqueous NaCl (20 mL), dried (Na₂SO₄) and concentrated on a rotary evaporator to yield 0.356 g (79%) of 1-indanone that was determined to be at least 96% pure by ¹H and ¹³C NMR spectroscopy and GC analysis.

In summary, this work demonstrates a new and useful method for deprotection of ketoximes. The advantages of this method include (1) the use of relatively non-toxic reagents that are insensitive to air and small amounts of moisture and (2) the use of a relatively non-toxic solvent system.

Acknowledgements

We gratefully acknowledge financial support from The National Science Foundation (NSF-RUI grant, CHE0078881). R.M. would also like to acknowledge The Camille and Henry Dreyfus foundation for a Henry Teacher Scholar Award.

References

1. (a) Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*, 3rd ed.; John Wiley and Sons: New York, 1999; (b) Hanson, J. R. *Protecting Groups in Organic Synthesis*, 1st ed.; Blackwell Science: Malden, MA, 1999; (c) Kocienski, P. J. *Protecting Groups*, 1st ed.; Georg Thieme Verlag: Stuttgart, 1994.
2. For some methods, see: *Comprehensive Organic Functional Group Transformations*, 1st ed.; Katritzky, A. R.; Meth-Cohn, O.; Rees, C. W., Eds.; Pergamon: New York, 1995; Vol. 3.
3. Bose, S. D.; Narsaiah, A. V. *Synth. Commun.* **1999**, *29*, 937.
4. Maloney, J. R.; Lyle, R. E. *Synthesis* **1978**, 212.
5. Barhate, N. B.; Gajare, A. S.; Waharkar, R. D.; Sudalai, A. *Tetrahedron Lett.* **1997**, *38*, 653.
6. Yadav, J. S.; Sasmal, P. K.; Chand, P. K. *Synth. Commun.* **1999**, *29*, 3667.
7. Demir, A. S.; Tanyeli, C.; Altinel, E. *Tetrahedron Lett.* **1997**, *38*, 7267.
8. Hajipour, A. R.; Mahboubghah, N. *OPPI BRIEFS* **1999**, *31*, 112.
9. Garrett, R. L. In *Designing Safer Chemicals*. American Chemical Society Symposium Series 640: Washington, DC, 1996, Chapter 1.
10. Ranu, B. C.; Sarkar, D. C. *J. Org. Chem.* **1988**, *53*, 878.
11. (a) Reglinski, J. In *Chemistry of Arsenic, Antimony and Bismuth*; Norman, N. C., Ed.; Blackie Academic and Professional: New York, 1998; pp. 403–440; (b) Marshall, J. A. *Chemtracts* **1997**, 1064–1075; (c) Suzuki, H.; Ikegami, T.; Matano, Y. *Synthesis* **1997**, 249; (d) Leonard, N. M.; Wieland, L. C.; Mohan, R. S. *Tetrahedron* **2002**, *58*, 8373.
12. Boruah, A.; Baruah, B.; Prajapati, D.; Sandhu, J. S. *Tetrahedron Lett.* **1997**, *38*, 4267.
13. (a) Hajipour, A. R.; Mallakpour, S. E.; Mohammadpoor-Baltork, I.; Adibi, H. *Monatshesfte fuer Chemie* **2003**, *134*, 45; (b) Hajipour, A. R.; Mallakpour, S. E.; Baltork, I. M.; Adibi, H. *Synth. Commun.* **2001**, *31*, 3401.
14. Nattier, B. A.; Eash, K. J.; Mohan, R. S. *Synthesis* **2001**, 1010.
15. *Preparation of bismuth triflate*: Labrouillère, M.; Le Roux, C.; Gaspard, H.; Laporterie, A.; Dubac, J.; Desmurs, J. R. *Tetrahedron Lett.* **1999**, *40*, 285. Bismuth triflate synthesized by this procedure is reported to be mainly the tetrahydrate. Recently, another procedure for the synthesis of bismuth triflate has been reported. Répichet, S.; Zwick, A.; Vendier, L.; Le Roux, C.; Dubac, J. *Tetrahedron Lett.* **2002**, *43*, 993.
16. Brezinski, B.; Grech, E.; Malarski, Z.; Sobozyk, L. *J. Chem. Soc., Perkin Trans. 2* **1991**, *2*, 857.
17. Juskowiak, M.; Krzyzanowski, P. *J. Prakt. Chemie.* **1989**, *331*, 870.